



Glowing products for science

Goat Anti-Rabbit IgG (H+L), F(ab')₂ Fragment

Goat anti-rabbit IgG (H+L), F(ab')₂ fragment secondary antibody labeled with our superior CF® Dyes and other labels



Product Description

This is a goat anti-rabbit IgG (H L) F(ab')₂ fragment secondary antibody labeled with our superior CF® Dyes and other labels.

- Available in 14 bright and photostable CF® Dyes, and HRP
- Biotin, APC, and R-PE conjugates also available
- Suitable for western, immunofluorescence, and immunohistology in FFPE tissues

Note: Conjugates of blue fluorescent dyes like CF®350 are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Call us : [800-304-5357](tel:800-304-5357)

Product attributes

Antibody type	Secondary
Clonality	Polyclonal
Host species	Goat
Antibody reactivity (target)	Rabbit IgG
Species reactivity	Rabbit
Cross adsorption	Not cross-adsorbed
Concentration	2 mg/mL
Antibody/conjugate formulation	Liquid: PBS/50% glycerol/2 mg/mL BSA/0.05% azide, Lyophilized: PBS/15 mg/mL BSA/20 mg/mL trehalose after reconstitution
Secondary/tag antibody applications	ELISA, Flow cytometry, IHC, IF (cells or tissue sections), Western blot
Product origin	Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.

Goat Anti-Rabbit IgG (H+L), F(ab')₂ Fragment

Conjugation	Ex/Em	Size	Catalog No.	Dye Features
CF@350 0.25 mL (500 ug)	347/448 nm 20146	50 uL (100 ug)	20146-1	CF@350 Features
CF@488A 0.25 mL (500 ug)	490/515 nm 20013	50 uL (100 ug)	20013-1	CF@488A Features
CF@505 0.25 mL (500 ug)	505/519 nm 20915-250uL	50 uL (100 ug)	20915-50uL	
CF@543 0.25 mL (500 ug)	541/560 nm 20330	50 uL (100 ug)	20330-1	CF@543 Features
CF@555 0.25 mL (500 ug)	555/565 nm 20035	50 uL (100 ug)	20035-1	CF@555 Features
CF@568 0.25 mL (500 ug)	562/583 nm 20099	50 uL (100 ug)	20099-1	CF@568 Features
CF@583R 0.25 mL (500 ug)	585/609 nm 20913-250uL	50 uL (100 ug)	20913-50uL	CF@583R Features
CF@594 0.25 mL (500 ug)	593/614 nm 20153	50 uL (100 ug)	20153-1	CF@594 Features
CF@633 0.25 mL (500 ug)	630/650 nm 20131	50 uL (100 ug)	20131-1	CF@633 Features
CF@640R 0.25 mL (500 ug)	642/662 nm 20087	50 uL (100 ug)	20087-1	CF@640R Features
CF@647 0.25 mL (500 ug)	650/665 nm 20045	50 uL (100 ug)	20045-1	CF@647 Features
CF@660C 0.25 mL (500 ug)	667/685 nm 20917-250uL	50 uL (100 ug)	20917-50uL	CF@660C Features
CF@680 0.25 mL (500 ug)	681/698 nm 20064	50 uL (100 ug)	20064-1	CF@680 Features
CF@750 0.25 mL (500 ug)	755/779 nm 20919-250uL	50 uL (100 ug)	20919-50uL	CF@750 Features
R-PE 500 uL (250 ug)	496, 546, 565/578 nm 20358-500uL	200 uL (100 ug)	20358-200uL	
APC 500 uL (250 ug)	650/660 nm 20415-500uL	100 uL (50 ug)	20415-100uL	
Biotin 0.25 mL (500 ug)	N/A 20188	50 uL (100 ug)	20188-1	

View our full selection of [Secondary Antibodies](#), or search our catalog using our [Antibody Finder](#). Alternatively, you can view our [secondary antibody product listings](#) with catalog numbers.

CF® Dyes offer exceptional brightness and photostability. For more information see our [CF® Dye technology page](#).

Storage and Handling

Liquid format: Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Liquid format antibodies contain 50% glycerol and will not freeze at -20°C.

Lyophilized format: Store at -20°C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Reconstitute antibodies in water using the indicated volumes below:

CF® Dye and biotin conjugates: add 0.5 mL dH₂O

HRP or DNP conjugates: add 1 mL dH₂O

Add the indicated volume of water directly to the vial containing the lyophilized antibody and mix gently to dissolve. Store reconstituted antibody at -20°C and protect from light. Aliquot to avoid repeated freeze/thaw cycles. Alternatively, an equal volume of glycerol can be mixed with the reconstituted antibody so that it will remain liquid at -20°C.

Optional: A preservative such as 0.05% sodium azide (final concentration) can be added to CF® Dye and biotin conjugates. Do not add sodium azide to HRP conjugates.

Note: Storage of the antibody for more than a day at final working dilution is not recommended.

CF is a registered trademark of Biotium, Inc.

References

Download a list of [CF® dye references](#).

This datasheet was generated on May 8, 2026 at 06:43:48 AM. Visit product page to check for updated information before use.
Product link: <https://biotium-woo.supremeclients.com/product/goat-anti-rabbit-igg-hlfab2-fragment/>