

Goat Anti-Human IgA (alpha chain)

Goat anti-human IgA (α chain) secondary antibody labeled with our superior CF® dyes.



Product attributes

Clonality	Polyclonal
Antibody type	Secondary, Anti-Human Immunoglobulin
Concentration	2 mg/mL
Antibody/conjugate formulation	Liquid: PBS/50% glycerol/2 mg/mL BSA/0.05% azide, Lyophilized: PBS/15 mg/mL BSA/20 mg/mL trehalose after reconstitution
Species reactivity	Human
Host species	Goat
Antibody reactivity (target)	Human IgA
Secondary/tag antibody applications	Flow cytometry, IHC, IF (cells or tissue sections), Western blot
Product origin	Product may contain either bovine serum albumin (BSA) from bovine serum (<i>Bos taurus</i>), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.

Product Description

This is a goat anti-human IgA (α chain) secondary antibody labeled with our bright and photostable CF® dyes. The antibodies react specifically with IgA heavy chains (α chains) and not with immunoglobulin light chains. To minimize cross-reactivity, the antibodies are cross-adsorbed against human IgG prior to conjugation.

- Cross-adsorbed for specific staining with minimal background
- Available in 3 bright and photostable CF® dyes
- Suitable for western, immunofluorescence, and immunohistology

Goat Anti-Human IgA (alpha chain)

Conjugation	Ex/Em	Size	Catalog No.	Dye Features
CF@488A 0.5 mL (1 mg) 1 mg (lyophilized)	490/515 nm 20428-500uL 20428-1mg	50 uL (100 ug)	20428-50uL	CF@488A Features
CF@594 0.5 mL (1 mg) 1 mg (lyophilized)	593/614 nm 20429-500uL 20429-1mg	50 uL (100 ug)	20429-50uL	CF@594 Features
CF@633 0.5 mL (1 mg) 1 mg (lyophilized)	630/650 nm 20427-500uL 20427-1mg	50 uL (100 ug)	20427-50uL	CF@633 Features

View our full selection of bright and specific [Secondary Antibodies](#), or search our catalog using our [Antibody Finder](#). Alternatively, you can view our [secondary antibody product listings](#) with catalog numbers.

CF® Dyes offer exceptional brightness and photostability. For more information see our [CF® Dye technology page](#).

Storage and Handling

Liquid format: Store at -20 °C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Liquid format antibodies contain 50% glycerol and will not freeze at -20 °C.

Lyophilized format: Store at -20 °C, protected from light. Product is stable for at least 6 months from date of receipt when stored as recommended. Reconstitute antibodies in water using the indicated volumes below:

CF® Dye and biotin conjugates: add 0.5 mL dH₂O

HRP or DNP conjugates: add 1 mL dH₂O

Add the indicated volume of water directly to the vial containing the lyophilized antibody and mix gently to dissolve. Store reconstituted antibody at -20 °C and protect from light. Aliquot to avoid repeated freeze/thaw cycles. Alternatively, an equal volume of glycerol can be mixed with the reconstituted antibody so that it will remain liquid at -20 °C.

Optional: A preservative such as 0.05% sodium azide (final concentration) can be added to CF® Dye and biotin conjugates. Do not add sodium azide to HRP conjugates.

Note: Storage of the antibody for more than a day at final working dilution is not recommended.

CF is a registered trademark of Biotium, Inc.

References

Download a list of [CF® dye references](#).

This datasheet was generated on May 9, 2026 at 07:20:57 PM. Visit product page to check for updated information before use.

Product link: <https://biotium-woo.supremecients.com/product/goat-anti-human-iga-alpha-chain/>