

## Biotin NTA

Biotin NTA, also known as BNTA (Biotin nitrilotriacetic acid, tripotassium salt) is used to detect histidine-tagged proteins (as little as 0.1 pmol) immobilized on nitrocellulose membranes.




### Product attributes

## Product Description

Biotin NTA, also known as BNTA (Biotin nitrilotriacetic acid, tripotassium salt) is used to detect histidine-tagged proteins (as little as 0.1 pmol) immobilized on nitrocellulose membranes. Biotin NTA can be loaded with Ni by combining it with a 3 fold molar excess of NiSO<sub>4</sub> at pH7.9 (1). Biotin NTA can be removed from the histidine-tagged protein at pH 4.8, allowing the blot be reanalyzed with another probe.

- White solid soluble in water
- Store at 4°C
- C<sub>20</sub>H<sub>36</sub>N<sub>4</sub>O<sub>8</sub>S
- MW: 488.56



## References

1. Anal Biochem 236, 101 (1996).
2. Anal Biochem 229, 119 (1995).

This datasheet was generated on May 10, 2026 at 11:16:48 AM. Visit product page to check for updated information before use.  
Product link: <https://biotium-woo.supremeclients.com/product/biotin-nta-biotin-nitrilotriacetic-acid-tripotassium-salt-also-called-bnta/>